DISCOVERY OF AJUGA CHAMAEPITYS (LAMIACEAE) NATURALIZED IN TEXAS

JEFFREY N. MINK

3229 Cole Ave. Waco, Texas 76707 jeffreynormanmink@gmail.com

JASON R. SINGHURST

Wildlife Diversity Program
Texas Parks and Wildlife Department
4200 Smith School Road
Austin, Texas 78744
jason.singhurst@tpwd.state.gov

WALTER C. HOLMES

Department of Biology
Baylor University
Waco, Texas 76798-7388
walter holmes@baylor.edu

ABSTRACT

Ajuga chamaepitys is reported as naturalized in a horse pasture along the eastern banks of White Oak Creek in Gillespie County of the Edwards Plateau. This is the second naturalized species of Ajuga known in Texas. The restricted distribution of this species suggests that it was established by adulterated forage transported with horse stock.

Ongoing floristic inventory in the Edwards Plateau of south-central Texas has resulted in the discovery of naturalized populations of *Ajuga chamaepitys* (L.) Schreb. This is the third state of the USA where this species has been documented (Virgina, Maryland, and Texas).

Voucher specimens: TEXAS. Gillespie Co.: Terra Linda Estates, ca. 15 km at 356 degrees from Kerrville (Kerr Co.), lat. 30°10'43.58", long. 99°08'55.62", ca. 595 m elev., 6 Apr 2015, W.C. Holmes, J.R. Singhurst & J.N. Mink 16134 (BAYLU); same general location, lat. 30°11'01.22", long. 99°08'41.63", same collectors and date, 16135 (BAYLU); same general location, lat. 30°11'18.68", long. 99°08'11.30", same collectors and date, 16130 (BAYLU). Figure 1.

The Gillespie County pasture, which is about 1.9 km long and varies from about 50 to 230 m in width, was densely and evenly covered with Ajuga. Associated vegetation within the pasture included Nassella leucotricha, Salvia farinacea, Marrubium vulgare, Glandularia bipinnatifida, Verbena halei, Ratibida columnifera, Lesquerella recurvata, Scutellaria drummondii, Arenaria sp., Evax sp., and Chaerophyllum tainturieri. Peripheral herbaceous vegetation and woody shrubs and trees included Desmodium paniculatum, Desmodium psilophyllum, Ulmus crassifolia, Juniperus ashei, Quercus buckleyi, Celtis laevigata, with Taxodium distichum and Platanus occidentalis along the littoral zone of White Oak Creek.

The genus Ajuga comprises 40-50 species of decumbent and erect annuals and perennials in North Africa, Mediterranean Europe, Great Britain, Israel, Asia, and Australia, distributed mostly in cooler regions (Cheifetz et al. 1999). Ajuga chamaepitys is native to central and southern Europe, the eastern Mediterranean region, and North Africa. None of the species are native to the Western Hemisphere (Diggs et al. 1999).

Until now, only Ajuga reptans L. was known from Texas (Cory & Parks 1937; Gould 1962; Correll & Johnston 1970; Hatch et al 1990; Johnston 1990; Jones et al. 1997; Turner et al. 2003; Hannick et al 2013). Ajuga chamaepitys was first reported in the USA in 1981 from Virginia (Harvill et al. 1981) and later from Maryland (Brown & Brown 1984). Three other species of Ajuga are known adventives or have escaped cultivation in North America (Radford et al. 1965; Strausbaugh & Core 1977; Kartesz 2015): (1) A. genevensis, in the northeastern USA and southeastern Canada (Gleason & Cronquist 1991; Brouillet et al. 2006; Kartesz 2015), (2) A. reptans, widespread throughout the USA except the Dakotas, Minnesota, Nebraska, Wyoming, Colorado, New Mexico, Nevada, California and Arizona (Gleason & Cronquist 1991; Brouillet et al. 2006; Kartesz 2015), and (3) A. pyramidalis, known only from Washington County, Mississippi (Bryson & Skojac 2011). The following key, adapted from Stace (1997), distinguishes the three species in the southern USA.

KEY TO AJUGA NATURALIZED IN SOUTHERN USA

- 1. Annuals; leaves dissected; plants neither stoloniferous nor rhizomatous; corollas yellow

 Ajuga chamaepitys

 1. Perennials; leaves subentire or serrate; plants stoloniferous or rhizomatous; corollas blue, pink or white.
 - 2. Plants rhizomatous; all bracts longer than flowers; stem puberulence continuous around stems

 Ajuga pyramidalis

 2. Plants stoloniferous; upper bracts shorter than flowers; upper part of stem puberulence discontinuous around stems (only on 2 opposite sides)

 Ajuga reptans

Ajuga chamaepitys in the Texas flora may remain localized, given its disjunct pattern of distribution in the USA (Virginia-Maryland and Texas), coupled with its known edaphic requirements of limestone, calcareous substrates, and calciferous soils (Lousley 1950; Stace 1997). Although Turrill (1948) described a large-scale ecological cline in the geographic distribution of the species from Asia Minor across Europe, this plant is designated by others as uncommon, decreasing or restricted in England (Lousley 1950; Stace 1997). Jim Scuddy, land manager of Terra Linda Estates, observes that the plant has been present in the Gillespie County horse pasture since the 1980s; no dispersal of A. chamaepitys has been observed outside of this area.

The establishment of Ajuga chamaepitys in the USA probably is connected to horseracing and horse recreational pursuits and the importation of horse stock from Europe and the Middle East (Weatherby 1791; Shouse 1908; Prior 1935; Wentworth 1938; Lightcap 1940) — seeds may have arrived via ruminant gut remnants or adulterated forage. Livestock forage as a vector for plant propagules is well known (Dewey 1897; Hillman & Henry 1928; Clines 2005) and interstate regulations related to weed-free hay are lax.

ACKNOWLEDGEMENTS

We thank Jim Scuddy, Ranch Manager of Terra Linda Estates, who brought the first sample of the species to Jim and Priscilla Stanley, naturalists and residents of Terra Linda Estates. Photos were then passed on to the authors, which led to plans for visiting the site. We also thank Jeff Forman, Wildlife Biologist at Mason Mountain Wildlife Management Area (MMWMA), who suggested that the plant might be *Ajuga chamaepitys* and Bill Carr of the University of Texas Herbarium for confirming the identification.



Figure 1. Ajuga chamaepitys, ground-pine or yellow bugle, in Gillespie Co., Texas, 6 April 2015. Photo by Jason Singhurst.

LITERATURE CITED

Brown, M.L. and R.G. Brown. 1984. Herbaceous Plants of Maryland. Port City Press, Baltimore, Maryland.

Brouillet, L., F. Coursol, and M. Favreau. 2006. VASCAN, the Database of Vascular Plants of Canada. Herbier Marie-Victorin, Institut de Recherche en Biologie Végétale, Univ. de Montréal.

Bryson, C.T. and D.A. Skojac, Jr. 2011. An annotated checklist of the vascular flora of Washington County, Mississippi. J. Bot. Res. Inst. Texas 5: 855–866.

Cheifetz, A., C. Double, L. Barnard, and D. Imwold (eds.). 1999. Botanica's Complete Annuals & Perennials. Laurel Glen Publishing, San Diego, California.

Clines, J. 2005. Preventing weed spread via contaminated hay and straw. Pp 4–6, <u>in</u> G. Skurka, (ed.). Proceedings of the California Invasive Plant Council Symposium. Volume 9. California Invasive Plant Council (Cal-IPC), Berkeley.

Cory, V.L. and H.B. Parks. 1937. Catalogue of the Flora of Texas. Texas Agric. Exp. Sta. 550, College Station.

Correll, D.S. and M.C. Johnston. 1970. Manual of the Vascular Plants of Texas. Texas Research Foundation, Renner.

Dewey, L.H. 1897. Migration of weeds. Pp. 263–286, in United States Department of Agriculture. Yearbook. Government Printing Office, Washington, D.C.

Diggs, G.M., B.L. Lipscomb, and R.J. O'Kennon. 1999. Illustrated Flora of North Central Texas. Sida Bot. Misc. 16. Bot. Res. Inst. of Texas, Fort Worth.

- Gleason, H.A. and A. Cronquist. 1991. Manual of Vascular Plants of the Northeastern United States and Adjacent Canada. New York Botanical Garden, Bronx, New York.
- Gould, F.W. 1962. Texas Plants—A Checklist and Ecological Summary. Texas Agricultural Experiment Station Publ. MP-585, Agricultural and Mechanical College of Texas, College Station.
- Hannick, V.C., J.N. Mink, J.R. Singhurst, and W.C. Holmes. 2013. Annotated checklist of the vascular flora of McLennan County, Texas. Phytoneuron 2013–29: 1–37.
- Harvill, A.M., C.E. Stevens, and D.M.E. Ware. 1981. Atlas of the Virginia flora, Part II. Dicotyledons. Virginia Botanical Associates, Farmville.
- Hatch, S.L., K.N. Gandhi, and L.E. Brown. 1990. Checklist of the Vascular Plants of Texas. Texas Agricultural Experiment Station Publ. MP-1655. Texas A&M University, College Station.
- Hillman, F.H. and H.H. Henry. 1928. The incidental seeds found in commercial seed of alfalfa and red clover. Proceedings of the International Seed Testing Association. October, No. 6: 1–20.
- Johnston, M.C. 1990. The Vascular Plants of Texas: A List, Updating the Manual of the Vascular Plants of Texas (ed. 2). Published by the author, Austin, Texas.
- Jones, S.D, J.K. Wipff, and P.M. Montgomery. 1997. Vascular Plants of Texas: A Comprehensive Checklist Including Synonymy, Bibliography, and Index. Univ. of Texas Press, Austin.
- Kartesz, J.T. 2015. Taxonomic Data Center. The Biota of North America Program (BONAP). Chapel Hill, North Carolina. http://www.bonap.net/tdc
- Lightcap, G.C. 1940. The American Thoroughbred. Iowa State Univ. Veterinarian: Vol. 2: Iss. 2, Article 1.
- Lousley, F.E. 1950. The New Naturalist—Wild Flowers of Chalk and Limestone. Collins, London.
- Prior, C.M. 1935. The Royal Studs of the Sixteenth and Seventeenth Centuries. Horse and Hound Publications Ltd., London.
- Radford, A.E., H.E. Ahles, and C.R. Bell. 1965. Atlas of the Vascular Flora of the Carolinas. Univ. of North Carolina, Dept. of Botany Techn. Bull. 165, Chapel Hill.
- Shouse, J. 1908. The American thoroughbred horse. J. Hered. 1: 92–106.
- Stace, C. 1997. New Flora of the British Isles (ed. 2). Cambridge Univ. Press, Cambridge, United Kingdom.
- Strausbaugh, P.D. and E.L. Core. 1977. Flora of West Virginia, (ed. 2 in 4 vols.). West Virginia Bulletin, Morgantown.
- Turner, B.L., H. Nichols, G. Denny, and O. Doron. 2003. Atlas of the Vascular Plants of Texas, Vol. I. Sida, Bot. Misc. 24. Botanical Res. Inst. of Texas, Fort Worth.
- Turrill, W.B. 1948. The New Naturalist British Plant Life. Collins, London.
- Weatherby, J. 1791. An Introduction To a General Stud Book. Weatherby and Sons, London.
- Wentworth, L. 1938. Thoroughbred Racing Stock. George Allen & Unwin Ltd., London.